

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

and these same fruit bodies were found to retain their vitality when dried, kept *in vacuo*, and at a temperature of liquid air for three weeks.—I. M. C.

Seedling anatomy.—In continuing his investigations of the seedling anatomy of Sympetalae, Lee¹⁰ has published an account of the Compositae, having examined about 50 species, well distributed through the tribes. The general conclusions are as follows: all seedlings are either diarch or tetrarch; variations in vascular anatomy occur not only in nearly related species, but in different individuals of the same species, the inference being that seedling anatomy "is of no value in questions of affinity"; the evolution of the vascular structures of seedlings is probably not an extremely slow process; tetrarchy and diarchy have probably been "interchanged" several times during the evolution of angiosperms; physiological factors are probably not sufficient to account for all the structures found in seedlings.—J. M. C.

The origin of Ascomycetes.—In a paper which reviews all the available data, approximately 100 papers being cited, Dodget discusses the relationships of the red algae and the Ascomycetes. It is a very useful summary of our knowledge of the reproductive structures of these two groups, as well as a clearly presented argument in favor of the view that the Ascomycetes are a monophyletic group and have been derived from the red algae. The reproductive structures of the two groups are compared in detail, and the interesting transitions shown by *Collema* and *Ascobolus* are described.—J. M. C.

Flora of Panama.—Standley¹² has issued the first of a series of papers preliminary to a flora of Panama. The present paper contains descriptions of some 40 new species from tropical America, which are distributed among 18 genera belonging to the Cyperaceae, Leguminosae, Gentianaceae, and Rubiaceae. Two new generic names are proposed, namely *Nothophlebia* and *Geocardia* (*Geophila* D. Don, not Berg.) of the Rubiaceae, and the following genera have been revised: *Sommera* (5), *Watsonamra* (11), and *Cobaea* (18).—J. M. Greenman.

Variation in Oenothera ovata.—Mrs. Brandegee¹³ has discovered that this Californian species of *Oenothera* has a remarkable range of variation. Apparently it is a plexus of "elementary species" quite as numerous as have been found in *O. Lamarckiana* and *O. biennis.*—J. M. C.

¹⁰ Lee, E., Observations on the seedling anatomy of certain Sympetalae. II. Compositae. Ann. Botany **28**:303–329. figs. 2. 1914.

¹¹ DODGE, B. O., The morphological relationships of the Florideae and the Ascomycetes. Bull. Torr. Bot. Club 41:157-202. figs. 13. 1914.

¹² STANDLEY, PAUL C., Studies of tropical American Phanerogams, no. 1. Contrib. U.S. Nat. Herb. 17:427-458, pls. 24-31. 1914.

¹³ Brandegee, Katharine L., Variation in *Oenothera ovata*. Univ. Calif. Publ. Bot. 6:41-50. pls. 8, 9. 1914.